

THE ROLE OF RUMEN CILIATE PROTOZOA IN RUMINAL FERMENTATION, DIGESTION AND ENTERIC METHANE PRODUCTION

Nguyen Hung Son and Nguyen Thi Duong Huyen

Faculty of Animal Science, Vietnam National University of Agriculture, Hanoi, Vietnam

Corresponding author: nhson@vnua.edu.vn. Tel: 0915064542

ABSTRACT

First described in 1843, rumen ciliate protozoa were considered to be important in ruminant nutrition and contributed up to 50% of the total microbial biomass in the rumen. Recent studies on the presence or absence of rumenciliate protozoa concluded that rumen protozoa are important, but not essential in the rumen ecosystem and to the well-being of host animals. Despite the fact that elimination of rumen protozoa (defaunation) negatively affects ruminal fermentation, plant cell wall digestion, defaunation results in an increase in the efficiency of bacterial protein synthesis and the rate of nitrogen flow to the duodenum, leading to increase average daily gain of host animals. Importantly, this increase in livestock productivity could occur alongside a reduction in enteric methane emissions.

Keywords: *ciliate, defaunation, microbial protein, methane*

INTRODUCTION

Rumen protozoa were first described in 1843 and considered to be important for the ruminant nutrition (Hungate, 1966). However, despite the fact that rumen protozoa contribute to as much as half the total microbial biomass in the rumen and actively participate in plant cell wall digestion (Williams and Coleman, 1992), elimination of rumen protozoa increases microbial protein supply to the host, leading to increase average daily gain by 11% and results in a 10.5 to 13% decrease in enteric methane production (Newbold et al., 2015; Eugène et al., 2004a). This paper aims at reviewing recent information on rumen ciliate protozoa and evaluating the role of protozoa on ruminal fermentation, digestion and enteric methane production.

Classification of rumen ciliate protozoa

Rumen protozoa are classified based on their cell morphology, and considered to be the simplest form of animal life, performing all the life processes as a eukaryotic cell (Dehority, 2003; Esteban et al., 2014). Further, the majority of protozoa identified in the rumen are ciliate species, with more than 100 species of rumen ciliate protozoa having been identified in two major sub-classes; being the entodiniomorphid (Table 1) and the holotrich ciliates (Table 2; Williams and Coleman, 1988, 1992). A few species of flagellate protozoa (Table 3) are also found in the rumen (Hungate, 1966; Williams and Coleman, 1992). However, the flagellates are easily confused with fungal zoospores (Dehority, 2003). In addition, flagellate protozoa are less numerous in terms of population density and have small body mass compared to the ciliates (Hungate, 1966; Clarke, 1977). Therefore, the flagellates are not well known and have not been the focus of attention in classifying or describing their activity and metabolism, leading to little information being available.

Table 1. Characteristics of some rumen entodiniomorphid protozoa (Williams and Coleman, 1988)

Genus	Dorsal cilia	Obvious skeletal plates	Macronucleus shape	Length (µm)	Width (µm)
Entodinium	0	0	Various	22-29	11-68
Eodinium	1 band ant.end [‡]	0	Rod-shaped	32-60	20-40
Diplodinium	1 band ant.end	0	Often bent rod	55-210	41-136
Eremoplastron	1 band ant.end	1 narrow	Often bent rod	45-500	21-260
Eudiplodinium	1 band ant.end	1 narrow	Hook shaped	105-198	56-120
Ostracodium	1 band ant.end	1 wide	Various	58-133	36-54
Polyplastron	1 band ant.end	2 narrow	Rod-shaped	123-205	98-123
Diploplastron	1 band ant.end	2 narrow close at post. End	Rod-shaped	88-120	47-65
Metadinium	1 band ant.end	2 narrow occ. Fused	Rod-shaped 2-3 lobes	110-288	61-165
Epidinium	1 band behind ant.end	3 variable width	Elongate	105-150	44-72
Enoploplastron	1 band ant.end	3 narrow close together	Elongate	60-140	32-90
Ophryoscolex	1 band round 3/4 of middle	3 variable width	Elongate	120-215	60-80
Epiplastron	1 band round 3/4 of middle	5 variables width	Elongate	90-140	41-60
Elytroplastron	1 band ant.end	3 narrow (2 on right 1 on left)	Elongate	110-160	67-97
Caloscolex	1 band round all middle	1 complex	Elongate	130-160	73-90
Opisthotrichum	1 band round 1/3 of middle	1 cylindrical	Elongate	60-80	21-28
Parentodium	0	0	Round	26-39	14-21

Note: [‡] anterior end

Table 2. Characteristics of some rumen holotrich ciliates (Williams and Coleman, 1992)

Species	Morphology	Size range, average (μm)	Length: width range, average	Macronucleus
Isotricha prostoma	Elongated ovoid to elipsoidal	80-200×50-120; 135×70	1.69-2.55; 2.03	Elongated
Isotricha intestinalis	Elongated ovoid to elipsoidal	90-200×45-150; 110×60	1.65-1.93; 1.76	Ovoid, 30×20
Dasytricha ruminantium	Ovoid	35-75×20-40; 57×27	1.70-2.70; 2.11	Elongated/ellipsoidal, 16-18×8-9
Dasytricha hukuokaensis	Ovoid	120-180×68-122; 151×95	1.47-1.76; 1.59	Ellipsoildal, 24-38×16-20; 31×18
Oligoisotricha bubali	Ovoid	12-20×8-15; 16×12	1.07-1.60; 1.30	Spherical-elliptical
Buetschlia parva	Ovoid	30-67×20-48; 55×35	1.58-2.38; 1.91	Spherical
Buetschlia neglecta	Ovoid	40-60×20-30	2.0	Spherical
Buetschlia lanceolate	Spear-shaped	48×20	2.4	Large
Buetschlia omnivore	Ovoid/spherical	Variable; 35-110×27-97		Elongated
Buetschlia nana	Ovoid	17-21×12-17; 19×15		spherical
Parabundleia ruminantium	Ovoid	37.5-50 × 27.5-32.5; 42.5 × 30.5	1.25-1.54	Elliptical, 16 μm long
Polymorphella bovis	Ovoid to bottle-like	26-37.5 × 20-26; 34× 22	1.30-1.80; 1.56	Subspherical, 2.5 μm long
Blepharoprosthium parvum	Pyriform	26-32 × 16-20; 29 × 18		Spherical
Blepharoconus krugerensis	Ovoid with anterior knoblike protuberence	30-65 × 21-60; 46 × 35	1.11-1.80; 1.34	Disc-shaped 7-15 × 4-8; 11 × 5.5 μm
Microcetus lappus	Ovoid/elongate	18-29 × 7.5-18; 23.6 × 13		Spherical

Table 3. Characteristics of some rumen flagellate protozoa (Williams and Coleman, 1992)

Species	Shape	Size	Number of flagella	Size of nuclear (μm)
<i>Chilomastix caprae</i>	Piriform	8.3×4.4	4	
<i>Monocercomonas ruminantium</i>	Piriform	4.8×4.1	4	1.8×1.6
<i>Monocercomonoide bovis</i>	Elliptical	5.4×2.8	4	1.6×1.4
<i>Monocercomonoide caprae</i>	Elliptical	9×6	4	Large
<i>Pentatrichomonas hominis</i>	Elliptical	7.5×5.6	5	2.5×2.0
<i>Tetratrichomonas buttrei</i>	Elliptical	5.3×4.8	4	2.0×1.7
<i>Trichomonas ruminantium</i>	Elliptical	12×10	3	

Anaerobic rumen ciliates are extremely abundant, ranging from 10^4 to 10^6 cells/mL of rumen liquor and are capable of engulfing bacteria and digesting plant materials such as cellulose and other structural carbohydrates (Finlay and Esteban, 2013; Esteban et al., 2014).

Rumen protozoa account for as much as half the total microbial biomass in the rumen and up to 50% of total fermentation products (Williams and Coleman, 1992; Newbold et al., 2015). They actively participate in the ruminant digestion process. The absence of rumen protozoa, therefore, results in modifying ruminal digestion of plant cell walls and starch which are considered to be two main sources of energy supply for ruminants (Jouany and Martin, 1997). The ruminal ecosystem and environment can be slightly altered by the absence of rumen protozoa with significant influence on bacterial activity, affecting the retention time of the digesta, the concentrations and proportion of ruminal VFA and ammonia (NH_3) concentration (Eugène et al., 2004a; Newbold et al., 2015) and therefore the supply of metabolites to the host, especially amino acids.

Plant cell wall digestion

Early studies on the role of rumen protozoa in ruminal digestion concluded that rumen protozoa did not digest plant cell components (Becker, 1929). In later years, enzymatic and microscopic evidence showed that cellulose was digested by entodiniomorphs (Hungate, 1966). Further, the rumen ciliate protozoa were confirmed in their ability to colonize and damage plant tissues in studies with scanning electron microscopic technology. *Epidinium crawleyi* was found to cause primary degradation of plant tissues (Bauchop and Clarke, 1976) as it attached and damaged areas of the stem. *Entodinium* spp. and *Ophryoscolex caudatum*, however, rarely attached themselves to the large tissue fragments, but to damaged tissues exposed through fractures (Orpin, 1984).

Rumen protozoa produce fibre-degrading enzymes and the cellulolytic enzymes produced by rumen protozoa are distinguished from those of bacterial and fungal origin. This was observed by the characterization of protozoal genes encoding cellulase enzymes (Jouany and Martin, 1997). Further, polysaccharidase activities were greater in animals with ciliate protozoa compared to animals without ciliate protozoa (Santra and Karim, 2002; Eugène et al., 2004b). However, effects of rumen protozoa on ruminal digestion are inconsistent in the literature. There are reports of decreased organic matter (OM) digestibility (Eugène et al., 2004a; Newbold et al., 2015), and neutral-detergent fibre (NDF) and acid-detergent fibre (ADF) digestibility (Newbold et al., 2015) in the absence of rumen protozoa, probably due to the loss of protozoal fibrolytic activity. The absence of rumen protozoa, in contrast, increases cellulolytic ruminococci populations (Mosoni et al., 2011). This increased cellulolytic ruminococci in the absence of rumen protozoa may compensate for the loss of protozoal fibrolytic activity, and therefore the digestibility of OM, NDF and ADF was not different between defaunated and faunated animals (Jouany et al., 1995). Zeitz et al. (2012) observed no effects of individual ciliate protozoa such as *Entodinium caudatum*, *Epidinium ecaudatum* and *Endiplodium maggii* on whole tract digestibility of OM, NDF or ADF and suggested that ciliate protozoa may not always improve plant cell wall digestion.

Carbohydrates and starch digestion

Both holotrich and entodiniomorph ciliate protozoa are believed to ferment carbohydrates to meet their energy requirement. Holotrich ciliates utilise soluble carbohydrates, while *Entodinium* spp. and *Epidinium* spp. preferably digest starch (Williams, 1989). De Smet et al. (1992) showed that the total protozoal population density was nearly double in a high concentrate diet compared to a high roughage diet, with holotrich ciliates increasing accordingly. *Entodinium* spp. and *Epidinium* spp. have their largest numbers with a high concentrate diet.

Wereszka and Michałowski (2012) found that *Diploplastron affine* possessed enzymes degrading starch and its derivatives. A protozoal cell extract for enzymatic studies found *Diploplastron affine* ciliate capable of digesting starch, released about 45 pmol VFA per protozoa per hour and utilised liberated energy for their energy requirement. The rate of starch degraded by *Diploplastron affine* is equivalent to $2.4 \pm 0.47 \mu\text{mol/L}$ glucose per mg protein per min and the degradation rate of maltose is approximately $0.05 \mu\text{mol/L}$ glucose per mg protein per min (Wereszka and Michałowski, 2012). The ciliate *Diploplastron affine* is also found to digest insoluble 1,3- β -glucans such as pachyman and 1,6- β -glucans such as pustulan as energy substrates (Belzecki et al., 2012).

Apart from being able to digest and utilise plant carbohydrates, rumen protozoa also ferment chitin of the rumen fungus. Morgavi et al. (1994) found *Piromyces* spp. strain OTS1 in monocultures or in the presence of rumen protozoa in vitro and reported that rumen protozoa adversely affect the growth of *Piromyces* spp. strain OTS1 and are able to digest fungal cell walls, resulting in 42% reduction of chitin which is a carbohydrate component of the fungal cell wall. *Diploplastron affine* and *Entodinium caudatum* are found to possess a chitinolytic

enzyme (Miltko et al., 2015b) and utilise chitin as a source of energy for ciliate metabolism (Miltko et al., 2015a).

Protein digestion and protozoal synthesis in the rumen

Ruminants with protozoa in the rumen (faunated) support a higher ruminal NH₃ concentration than do animals with rumen protozoa removed (defaunated), indicating that rumen protozoa degrade dietary proteins (Jouany, 1996) and engulf bacteria for their amino acid requirement (Coleman, 1989; Esteban et al., 2014). Ueda et al. (1975) who incubated the isotrich ciliates with soluble casein found that peptide-nitrogen and amino-nitrogen produced by the isotrich reached its highest level between 3 and 15 hours of incubation, accounting for 47% and 58% of non-protein nitrogen (NPN), respectively. Incubation of the ophryoscolecoid ciliates with insoluble casein showed a peak of peptide-nitrogen at 3 hours, which accounted for 36% of NPN while amino-nitrogen increased linearly and accounted for 47% of NPN at 15 hours of incubation (Ueda et al., 1975).

Protozoal nitrogen (N) can account for 53% of the total microbial N in the bovine rumen (Michałowski, 1979), which is about 24 to 46 g N (Leng et al., 1981). However, rumen protozoa contribute only 20% of the total microbial N entering the duodenum (Jouany et al., 1988). The smaller protozoal biomass in the duodenum of the ruminants could be due to 65% to 74% of protozoa lysing and being degraded in the rumen (Leng, 1982; Ffoulkes and Leng, 1988), suggesting only 24 to 35% of protozoa enter the lower digestive tract. The relatively high numbers of rumen protozoa that complete their life span in the rumen (Leng, 1982) and are retained within the omasum of ruminants (Czerkawski, 1987; Nguyen and Hegarty, 2019) mean rumen protozoa contribute a small proportion of the total microbial protein supply. The principal detrimental effect of rumen protozoa, therefore, may be competition for substrate with bacteria and engulfment and digestion of bacteria by protozoa, leading to decreased bacterial biomass and flow of protein in the duodenum (Leng, 1982).

Ruminal lipid metabolism

The role of rumen protozoa in bio-hydrogenation is not well defined and understood (Williams and Coleman, 1992), although they contribute significantly to flow of unsaturated fatty acids to the duodenum (Newbold et al., 2015; Yáñez-Ruiz et al., 2006). Yáñez-Ruiz et al. (2006) reported rumen protozoa accounted for between 30-40% of conjugated linoleic acid (CLA) and 40% of vaccenic acid (VA) leaving the rumen. Mixed protozoa from the sheep rumen contain at least two to three times more unsaturated fatty acids, including CLA and VA, than do bacteria. Different species have different composition, with larger fibrolytic species such as *Epidinium ecaudatum caudatum* containing more than ten times more CLA and VA than some small species, including *Entodinium nanellum* (Devillard et al., 2006). This high level of polyunsaturated fatty acids in protozoal cells is a consequence of ingestion and/or engulfment of chloroplasts (Huws et al., 2009) and this chloroplasts uptake is specifically found in entodiniomorphids (Huws et al., 2012). Rumen protozoa, therefore appear to increase the duodenal flow of mono or polyunsaturated fatty acids by protecting chloroplasts unsaturated fatty acids from rumen bio-hydrogenation.

Factors affecting protozoal population densities in the rumen

Rumen ciliate protozoa represent approximately 10^4 - 10^6 cells/mL of rumen contents (Dehority 2003; Esteban et al., 2014), but the concentration of rumen protozoa varies among animals and is dependent on many factors such as ruminant species, geographical location (Akbar et al., 2009), diet (Whitelaw et al., 1984), frequency of feeding and rumen pH (Clarke, 1977).

Diet composition

Ruminants fed highly digestible diets often show the largest populations of rumen protozoa (Hungate, 1966), while small populations of rumen protozoa are found in animals on low quality roughage diets (Abe et al., 1973). De Smet et al. (1992) fed sheep low and high concentrate diets containing 4.3% or 17.3% starch respectively, observing total protozoal population was nearly two-fold higher in the high concentrate diet. Rumen protozoa are able to reduce the rate of fermentation, contributing to the maintenance of a stable ruminal ecosystem when a high concentration of grain is suddenly introduced in the diet (Mackie et al., 1978). However, the protozoa are significantly affected by the environment's acidity or alkalinity, with the protozoa unable to survive if rumen pH is above 7.8 or below 5.0 (Clarke, 1977). Mackie et al. (1978) also found that the protozoal population decreased by 50-80% as rumen pH fell below 5.4. Cellulolytic ciliates almost disappeared when cattle were fed barley only (Kudo et al., 1990) and steers became protozoa-free for a period of a few weeks by ad libitum feeding of barley (Whitelaw et al., 1984).

Dietary fatty acid supplement

Capric acid (C10:0), lauric acid (C12:0) and myristic acid (C14:0) show strong protozoal toxicity and are useful rumen defaunating agents (Matsumoto et al., 1991). Matsumoto et al. (1991) observed that rumen protozoa, except Entodinium spp. were undetectable after 3 days of feeding 30 g of hydrated coconut oil (CO) containing 52% lauric acid. Feeding 250g of refined CO to beef heifers reduced rumen protozoal population by 62% (Jordan et al., 2006) and protozoal populations in beef heifers were decreased by 63% and 80% by 300 g/d CO after 45 and 75 days, respectively (Lovett et al., 2003). Machmüller (2006) reported a reduction in rumen protozoa by 88 and 97% when feeding sheep with 3.5 and 7% CO, respectively. This suppressive effect of CO on rumen protozoa persisted 5 weeks after finishing feeding sheep with CO (Sutton et al., 1983). Rumen protozoa were reduced to half of the original population by cottonseed, with holotrich and cellulolytic protozoa apparently lost from the rumen of sheep and only Entodinium spp. remained (Dayani et al., 2007).

Frequency of feeding

The concentration of protozoa in the rumen liquor varies according to the daily feeding regime, reaching a maximum before feeding and decreasing by approximately 60-80% from 4 to 12 hours after feeding (Michalowski and Muszyński, 1978). More specifically, the holotrich population in the fluid decreases for a period of 12 to 20 hours after feeding and the population returns to its original numbers within 4 to 6 hours pre-feeding, while the entodiniomorphid population in the fluid decreases for up to 16 hours after feeding and then

increases to the pre-feeding numbers (Williams, 1986). The increase of the holotrich population is mainly caused by the increase in dasytrichs while the isotrich population remains relatively low (Clarke, 1965). The highest concentration of rumen protozoa occurs when the animal is fed three or four meals per day rather than once (Bonhomme, 1990).

Defaunating the rumen

Rumen protozoa are important, but not essential in the rumen ecosystem and to the well-being of host animals (Williams and Coleman, 1992; Newbold et al., 2015). Elimination of rumen protozoa (defaunation) has led to reported increases in growth rate and liveweight gain of ruminants (Bird and Leng, 1978; Bird et al., 1979; Eugène et al., 2004a; Newbold et al., 2015) especially when the feed is deficient in protein relative to energy content (Nguyen et al., 2015). In addition, rumen protozoa are significant hydrogen (H₂) producers and synthesise mainly acetate and butyrate rather than propionate (Williams and Coleman, 1992). Defaunation is therefore expected to induce a greater proportion of propionate in the ruminal VFA (Eugène et al., 2004a), but this phenomenon is not always observed (Williams and Coleman, 1992; Newbold et al., 2015). The reduced CH₄ emissions caused by defaunation also reported by several authors (Whitelaw et al., 1984; Hegarty, 1999; McAllister and Newbold, 2008; Newbold et al., 2015) may reflect reduced H₂ availability by removing endosymbiotic methanogens (Finlay et al., 1994; Tokura et al., 1997; Finlay and Esteban, 2013).

Effect of defaunation on extent of ruminal fermentation

An in vitro study by Yoder et al. (1966) reported cellulose digestion by rumen protozoa (7%), by bacteria (40%) and by protozoa and bacteria combined (exceeded 60%), showing a beneficial effect of rumen protozoa on cellulose digestion. Bauchop and Clarke (1976) observed rumen ciliate protozoa contribute to fibre digestion as they are capable of colonizing and damaging plant tissues. Cellulolytic, polysaccharide depolymerases and glycoside hydrolase enzymes produced by protozoa are significant contributors to cellulose and hemicellulose fermentation (Coleman, 1989). Therefore, the absence of rumen protozoa can lead to a 5-15% reduction in carbohydrate digestion of plant cell walls (Jouany et al., 1988).

Removing protozoa reduces the rumen digestibility of fibre components of the diet (Newbold et al., 2015). Ruminal digestion of NDF and ADF were reduced by 31% and 22% respectively by defaunation of sheep when fed a low soluble N diet (Ushida and Jouany, 1990). Defaunation also reduced degradation of a mainly chopped hay diet by up to 18% in sacco (De Smet et al., 1992). The absence of rumen protozoa did not affect rumen digestibility in lambs offered a diet with a high protein/energy ratio, but reduced total tract digestibility of OM (10%) and NDF (7%; Eugène et al., 2010). In addition, Ushida and Jouany (1990) found that defaunated ruminants require an increased supply of NPN in order to maintain fibrolytic activity in the rumen compared to faunated animals. Although fibre digestion is moderately suppressed by defaunation, improving protein supply is far more important in growing animals with high protein demand and when protein is a limiting factor in the diet (De Smet et al., 1992).

Effect of defaunation on animal productivity

A positive effect of defaunation on ruminants is the increased rumen bacterial biomass and passage of ruminal undegraded protein from the diet (Jouany, 1996). Duodenal N and duodenal CP/kg DMI outflow significantly increased after defaunation (Eugène et al., 2004a; Newbold et al., 2015), indicating an increase in the efficiency of microbial protein synthesis and leading to an average increased daily gain of 11%.

In pen-feeding studies, defaunated lambs showed 18% faster growth rate and greater wool growth and wool fibre diameter over faunated or refaunated lambs offered a 50:50 concentrate and roughage ration (Santra et al., 2007). Birth weight of lambs born from defaunated ewes was 13% heavier than from faunated ewes on single-born lamb and pre-weaning growth rates were 10% and 14% heavier in lambs reared free of ciliate protozoa for both single and twin-born lambs, respectively (Hegarty et al., 2008). On high energy and low protein diets, defaunated cattle grew at a 43% faster rate than faunated cattle on the same intake (Bird and Leng, 1978) and lambs without rumen protozoa showed significantly increased growth rates and efficiency of utilisation of feed when fed a low level of protein. Wool growth increased by 50% compared to faunated animals that were fed a low protein diet (Bird et al., 1979).

In grazing studies, Bird and Leng (1984) observed a greater rate of body weight gain (23%) and wool growth (19%) in defaunated compared to faunated lambs grazed on a green oats pasture. Protozoa-free lambs born from defaunated ewes were significantly (4-8%) heavier than were lambs born from faunated ewes measured from 2 months of age to 5 months of age and wool growth was also greater in protozoa-free lambs grazed on fescue dominant pastures (Hegarty et al., 2000).

Effect of defaunation on enteric methane production

As stated earlier, ciliate protozoa are significant producers of H₂ and produce acetic and butyric acids rather than propionic acid (Williams and Coleman, 1992). Defaunation is generally associated with fermentation shifting to a greater proportion of propionic acid, therefore reducing the amount of CH₄ produced (Eugène et al., 2004a).

The methanogens existing as endo- and ecto-symbionts with ciliate protozoa (Finlay et al., 1994; Tokura et al., 1997; Finlay and Esteban, 2013) have been estimated to account for 37% of ruminal methane production (Finlay et al., 1994). The proportion of methanogens in the total bacterial population was lower in protozoa-free lambs, with 26% lower CH₄ emissions compared to faunated lambs (McAllister and Newbold, 2008). While the archaeal community of methanogens in liquid and solid rumen contents were similar in faunated wethers, a lower proportion of methanogens occurred in the liquid phase with defaunation (Morgavi et al., 2012). However, Mosoni et al. (2011) while observing a 20% reduction in CH₄ emissions in short-term (10 week) and long-term (2 year) defaunated sheep, found methanogens per gram of DM of rumen content increased with defaunation while the diversity of the dominant methanogenic community was not changed. Therefore, it may not be reasonable to attribute the reduced CH₄ production from defaunation to a loss of methanogens (Morgavi et al., 2012).

The presence of protozoa did not change enteric CH₄ production in lambs raised with/without protozoa from birth (Hegarty et al., 2008) or from 10 to 25 weeks after chemical defaunation (Bird et al., 2008). Defaunation was associated with a reduced number of methanogens in rumen fluid, but did not reduce CH₄ production (Morgavi et al., 2012; Kumar et al., 2013). This could be explained as defaunation induces changes in bacterial or fungal populations (Eugène et al., 2004a) and the absence of protozoa in the rumen leads to changes in the methanogen community (Morgavi et al., 2012).

Ruminal acetogens were found able to grow on CO₂ and H₂, and produce acetate, but reductive acetogenesis was not likely to be occurring because of lower H₂ affinity, making reductive acetogenesis unable to compete with methanogens in the rumen (Joblin, 1999). In normal fermentation, methanogens reduce H₂ to a low level in which reductive acetogenesis is below detectable levels (Ungerfeld, 2015), but if pyruvate-derived acetate is produced when methanogenesis is inhibited, H₂ may accumulate and stimulate reductive acetogenesis (Ungerfeld, 2013). Reductive acetogens established in the rumen lacking methanogens can replace methanogens as a sink for H₂ in the rumen (Fonty et al., 2007). The reduced CH₄ emissions from defaunated animals associated with a rise in acetate proportion is a desirable condition in the rumen where reductive acetogens may be occurring.

CONCLUSION

Rumen ciliate protozoa are important in the ruminant nutrition, but they are not essential in the rumen ecosystem and to the well-being of host animals. Though elimination of rumen ciliate protozoa slightly decreases total VFA and plant cell wall digestion, defaunation largely impacts on the reduced NH₃ concentration, CH₄ production in the rumen and increased microbial protein supply to the hosts.

REFERENCES

- Abe, M., Shibui, T., Iriki, T. and Kumeno, F. 1973 Relationship between diet and protozoal population in the rumen. *British Journal of Nutrition* 29, pp. 197-202.
- Akbar, T., Ali, M. S., Golamreza, Z., Maghsoud, B. and Adel, A. 2009. The study of diversity of ciliate protozoa in Ghizel sheep fed in pasture and nourished by dried grape by-product. *American Journal of Animal and Veterinary Sciences* 4, pp. 37-41.
- Bauchop, T. and Clarke, R. T. 1976. Attachment of the ciliate *Epidinium* Crawley to plant fragments in the sheep rumen. *Applied and Environmental Microbiology* 32, pp. 417-422.
- Becker, E. R. 1929. Methods of retendering the rumen and reticulum of ruminants free from their normal infusorian fauna. *Proceeding of the National Academy of Sciences. U.S.A* 15, pp. 435-438. doi: 10.1073/pnas.15.5.435
- Belzecki, G., Miltko, R. and Michalowski, T. 2012. Ability of rumen protozoa *Diploplastron* affine to utilize β-glucans. *Folia Microbiol* 57, pp. 259-262.
- Bird, S. G. and Leng, R. A. 1978. The effects of defaunation of the rumen on the growth of cattle on low-protein high-energy diets. *British Journal of Nutrition* 40, pp. 163-167. doi:10.1079/bjn19780108
- Bird, S. G. and Leng, R. A. 1984. Absence of protozoa in the rumen and production of grazing sheep. Available at. <http://livestocklibrary.com.au/handle/1234/7475?show=full>

- Bird, S. H. and Light, M. N. 2013. A protocol for the removal of protozoa (defaunation) from the rumen of cattle. In *Recent Advances in Animal Nutrition-Australia*. School of Environmental and Rural Science, University of New England, Armidale NSW 2351, Australia'. (Ed. PB, Cronje) pp. 1.
- Bird, S. H., Hegarty, R. S. and Woodgate, R. 2008 Persistence of defaunation effects on digestion and methane production in ewes. *Australian Journal of Experimental Agriculture* 48, pp. 152-155. doi: 10.1071/EA07298
- Bird, S. H., Hill, M. K. and Leng, R. A. 1979. The effects of defaunation of the rumen on the growth of lambs on low-protein-high-energy diets. *British Journal of Nutrition* 42, pp. 81-87. doi: <https://doi.org/10.1079/BJN19790091>
- Bonhomme, A. 1990. Rumen ciliates: their metabolism and relationships with bacteria and their hosts. *Animal Feed Science and Technology* 30, pp. 203-266.
- Clarke, R. T. J. 1965. Diurnal variation in the numbers of rumen ciliate protozoa in cattle. *New Zealand Journal of Agricultural Research* 8, pp. 1-9.
- Clarke, R. T. J. 1977. Protozoa in the rumen ecosystem. In *'Microbial ecology of the gut.'* (Eds RTJ, Clarke, T, Bauchop) pp. 251-275. (Academic Press: New York).
- Coleman, G. S. 1989. Protozoal-Bacterial interaction in the rumen. In *'The roles of protozoa and fungi in ruminant digestion.'* (Eds JV, Nolan, RA, Leng, DI, Demeyer) (Penambul books: Armidale, N.S.W 2351, Australia).
- Czerkawski, J. W. 1987. Reassessment of the contribution of protozoa to the microbial protein supply to the host ruminant animal. *Journal of Theoretical Biology* 126, pp. 335-341.
- Dayani, O., Ghorbani, G. R., Alikhani, M., Rahmani, H. R. and Mir, P. S. 2007. Effects of dietary whole cottonseed and crude protein level on rumen protozoal population and fermentation parameters. *Small Ruminant Research* 69, pp. 36-45.
- De Smet, S., Demeyer, D. I. and van Nevel, C. J. 1992. Effect of defaunation and hay: concentrate ratio on fermentation, fibre digestion and passage in the rumen of sheep. *Animal Feed Science and Technology* 37, pp. 333-344.
- Dehority, B. A. 2003. *Rumen microbiology*. (Nottingham University Press: Thrumpton, Nottingham, UK).
- Devillard, E., McIntosh, F. M., Newbold, C. J. and Wallace, R. J. 2006. Rumen ciliate protozoa contain high concentrations of conjugated linoleic acids and vaccenic acid, yet do not hydrogenate linoleic acid or desaturate stearic acid. *British Journal of Nutrition* 96, pp. 697-704.
- Esteban, G. F., Finlay, B. J. and Warren, A. 2014. Free-Living Protozoa. In *'Ecology and General Biology.'* (Eds JH, Thorp, DC, Rogers) Vol. 1, pp. 113-132. (Academic Press: Boston).
- Eugène, M., Archimède, H. and Sauvant, D. 2004a. Quantitative meta-analysis on the effects of defaunation of the rumen on growth, intake and digestion in ruminants. *Livestock Production Science* 85, pp. 81-97. [https://doi.org/10.1016/S0301-6226\(03\)00117-9](https://doi.org/10.1016/S0301-6226(03)00117-9)
- Eugène, M., Archimède, H., Weisbecker, J. H., Periacarpin, F. and Saminadin, G. 2004b. Effects of defaunation on digestion and growth, in sheep receiving a mixed diet (fresh *Digitaria decumbens* grass and concentrate) at four protein to energy ratios. *Animal Research, EDP Sciences* 53, pp. 111-125.
- Eugène, M., Sauvant, D., Weisbecker, J. L., Archimède, H. 2010. Effects of defaunation on digestion of fresh *Digitaria decumbens* grass and growth of lambs. *Animal* 4, pp. 439-445.
- Ffoulkes, D. and Leng, R. A. 1988. Dynamics of protozoa in the rumen of cattle. *British Journal of Nutrition* 59, pp. 429-436.
- Finlay, B. J. and Esteban, G. F. 2013. Protozoa. In *'Encyclopedia of Biodiversity.'* (Ed. BJ, Finlay) pp. 286-297. (Academic Press: Waltham).
- Finlay, B. J., Esteban, G., Clarke, K. J., Williams, A. G., Embley, T. M. and Hirt, R. P. 1994. Some rumen ciliates have endosymbiotic methanogens. *FEMS Microbiology Letters* 117, pp. 157-161. <https://www.sciencedirect.com/science/article/abs/pii/0378109794901880>

- Fonty, G., Joblin, K., Chavarot, M., Roux, R., Naylor, G. and Michallon, F. 2007. Establishment and development of ruminal hydrogenotrophs in methanogen-free lambs. *Applied and Environmental Microbiology* 73, pp. 6391-6403. http://www.uky.edu/Ag/AnimalSciences/instruction/asc684/PDF/AppEnvMicrobiol73_6391.pdf
- Hegarty, R. S. 1999. Reducing rumen methane emissions through elimination of rumen protozoa. *Australian Journal of Agricultural Research* 50, pp. 1321-1327. <https://doi.org/10.1071/AR99008>
- Hegarty, R. S., Bird, S. H., Vanselow, B. A. and Woodgate, R. 2008. Effects of the absence of protozoa from birth or from weaning on the growth and methane production of lambs. *British Journal of Nutrition* 100, 1220-7. doi: 10.1017/S0007114508981435.
- Hegarty, R. S., Shands, C., Harris, C. and Nolan, J. V. 2000. Productivity and pasture intake of defaunated crossbred sheep flocks. *Australian Journal of Experimental Agriculture* 40, pp. 655-662. doi: 10.1071/EA99091
- Hungate, R. E. 1966. *The rumen and its microbes*. (Academic Press: New York).
- Huws, S. A., Kim, E. J., Kingston-Smith, A. H., Lee, M. R., Muetzel, S. M., Cookson, A. R., Newbold, C. J., Wallace, R. J., Scollan, N. D. 2009. Rumen protozoa are rich in polyunsaturated fatty acids due to the ingestion of chloroplasts. *FEMS Microbiology Ecology* 69, pp. 461-471.
- Huws, S. A., Lee, M. R., Kingston-Smith, A. H., Kim, E. J., Scott, M. B., Tweed, J. K., Scollan, N. D. 2012. Ruminal protozoal contribution to the duodenal flow of fatty acids following feeding of steers on forages differing in chloroplast content. *British Journal of Nutrition* 108, pp. 2207-2214.
- Joblin, K. N. 1999. Ruminal acetogens and their potential to lower ruminant methane emissions. *Australian Journal of Agricultural Research* 50, pp. 1307-1314.
- Jordan, E., Lovett, D. K., Monahan, F. J., Callan, J., Flynn, B. and O'Mara, F. P. 2006. Effect of refined coconut oil or copra meal on methane output and on intake and performance of beef heifers. *Journal of Animal Science* 84, pp. 162-170. doi:10.2527/2006.841162x
- Jouany, J. P. 1996. Effect of rumen protozoa on nitrogen utilization by ruminants. *The Journal of Nutrition* 126, pp. 12-1335. doi: 10.1093/jn/126.suppl_4.1335s
- Jouany, J. P., Demeyer, D. I. and Grain, J. 1988. Effect of defaunating the rumen. *Animal Feed Science and Technology* 21, pp. 229-265. doi:10.1080/17450398509425190
- Jouany, J. P. and Martin, C. 1997. Effect of protozoa in plant cell wall and starch digestion in the rumen. In 'Rumen microbes and digestive physiology in ruminants.' (Eds. R. Onodera, H. Itabashi, K. Ushida, H. Yaakub, Y. Sasaki), pp. 11-24. (Japan Scientific Societies Press: Tokyo/S.Karger AG).
- Jouany, J. P., Senaud, J., Toillon, S., Ben Salah, M., Bohatier, J. and Prensier, G. 1995. Effect of ruminal inoculation of *Isotricha* alone or a mixed B-type fauna in a defaunated rumen on the digestion of a hay-maize diet (70:30) in sheep. *Reproduction, Nutrition, Development* 35, pp. 11-25.
- Kudo, H., Cheng, K. J., Imai, S., Han, S. S. and Costerton, J.W. 1990. Effects of feed on the composition of the rumen ciliate protozoal population in cattle and its relationship to cellulolytic ciliate protozoa. *Animal Feed Science and Technology* 29, pp. 159-169.
- Kumar, S., Dagar, S. S., Puniya, A. K. and Upadhyay, R. C. 2013. Changes in methane emission, rumen fermentation in response to diet and microbial interactions. *Research in Veterinary Science* 94, pp. 263-268. <https://doi.org/10.1016/j.rvsc.2012.09.007>
- Leng, R. A. 1982. Dynamics of protozoa in the rumen of sheep. *British Journal of Nutrition* 48, pp. 399-415.
- Leng, R. A., Gill, M., Kempton, T. J., Rowe, J. B., Nolan, J. V., Stachiw, S. J. and Preston, T. R. 1981. Kinetics of large ciliate protozoa in the rumen of cattle given sugar cane diets. *British Journal of Nutrition* 46, pp. 371-384.
- Lovett, D., Lovell, S., Stack, L., Callan, J., Finlay, M., Conolly, J. and O'Mara, F. P. 2003. Effect of forage/concentrate ration and dietary coconut oil level on methane output and performance of finishing beef heifers. *Livestock Production Science* 84, pp. 135-146. <https://doi.org/10.1016/j.livprodsci.2003.09.010>

- Machmüller, A. 2006. Medium-chain fatty acids and their potential to reduce methanogenesis in domestic ruminants. *Agriculture, Ecosystems & Environment* 112, pp. 107-114. <https://doi.org/10.1016/j.agee.2005.08.010>
- Mackie, R. I., C, F. M., Gilchrista, A. M., Robbertsa, P. E. and Helen, M. S. 1978. Microbiological and chemical changes in the rumen during the stepwise adaptation of sheep to high concentrate diets. *Journal of Agricultural Science*. Summary 90, pp. 241-254.
- Matsumoto, M., Kobayashi, T., Takenaka, A. and Itabashi, H. 1991. Defaunation effects of medium-chain fatty acids and their derivatives on goat rumen protozoa. *The Journal of General and Applied Microbiology* 37, pp. 439-445. doi: 10.2323/jgam.37.439
- McAllister, T. A. and Newbold, C. J. 2008. Redirecting rumen fermentation to reduce methanogenesis. *Australian Journal of Experimental Agriculture* 48, pp. 7-13. doi:10.1071/EA07218
- Michałowski, T. 1979. Effect of feeding frequency on the diurnal changes in microbial protein, volatile fatty acids and ammonia contents of the rumen of sheep. *The Journal of Agricultural Science* 93, pp. 67-70.
- Miltko, R., Kowalik, B., Michałowski, T. and Bełżecki, G. 2015a. Chitin as a source of energy for rumen ciliates. *Journal of Animal and Feed Sciences* 24, pp. 203-207.
- Miltko, R., Pietrzak, M., Bełżecki, G., Wereszka, K., Michałowski, T. and Hackstein, J. H. P. 2015b. Isolation and in vitro cultivation of the fibrolytic rumen ciliate *Eremoplastron* (*Eudiplodinium*) *dilobum*. *European Journal of Protistology* 51, pp. 109-117.
- Morgavi, D. P., Martin, C., Jouany, J. P. and Ranilla, M. 2012. Rumen protozoa and methanogenesis: not a simple cause-effect relationship. *The British Journal of Nutrition* 107, pp. 388-97. doi:10.1017/S0007114511002935
- Morgavi, D. P., Sakurada, M., Mizokami, M., Tomita, Y. and Onodera, R. 1994. Effects of ruminal protozoa on cellulose degradation and the growth of an anaerobic ruminal fungus, *Piromyces* sp. strain OTS1, in vitro. *Applied and Environmental Microbiology* 60, pp. 3718-3723.
- Mosoni, P., Martin, C., Forano, E. and Morgavi, D. P. 2011. Long-term defaunation increases the abundance of cellulolytic ruminococci and methanogens but does not affect the bacterial and methanogen diversity in the rumen of sheep. *Journal of Animal Science* 89, pp. 783-791. doi: 10.2527/jas.2010-2947
- Newbold, C. J., de la Fuente, G., Belanche, A., Ramos-Morales, E. and McEwan, N. 2015. The role of ciliate protozoa in the rumen. *Frontiers in Microbiology* 6, pp. 1-14. <https://doi.org/10.3389/fmicb.2015.01313>
- Nguyen, S. H. and Roger Stephen Hegarty. 2019. Distribution of ciliate protozoal populations in the rumen, reticulum, and omasum of Angus heifers offered lucerne cereal mix. *Livestock Research for Rural Development*. 31(9). <http://www.lrrd.org/lrrd31/9/nhson31146.html>
- Nguyen, S. H., Barnett, M. and Hegarty, R. S. 2015. Use of dietary nitrate to increase productivity and reduce methane production of defaunated and faunated lambs consuming protein deficient chaff. *Animal Production Science*. 56(3), pp. 290-297. doi: 10.1071/AN15525
- Orpin, C. G. 1984. The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls. *Animal Feed Science and Technology* 10, pp. 121-143. [https://doi.org/10.1016/0377-8401\(84\)90003-8](https://doi.org/10.1016/0377-8401(84)90003-8)
- Santra, A. and Karim, S. A. 2002. Influence of ciliate protozoa on biochemical changes and hydrolytic enzyme profile in the rumen ecosystem. *Journal of Applied Microbiology* 92, pp. 801-811. <https://sfamjournals.onlinelibrary.wiley.com/doi/pdf/10.1046/j.1365-2672.2002.01583.x>
- Santra, A., Parthasarathy, S. and Karim, S. A. 2007. Effect of defaunation on body conformation changes, wool yield and fibre characteristics of growing lambs in a hot semiarid environment. *Australian Journal of Experimental Agriculture* 47, pp. 789-795. doi: 10.1071/EA05106
- Sutton, J. D., Knight, R., McAllan, A. B. and Smith, R. H. 1983. Digestion and synthesis in the rumen of sheep given diets supplemented with free and protected oils. *British Journal of Nutrition* 49, pp. 419-432.
- Tokura, M., Ushida, K., Miyazaki, K. and Kojima, Y. 1997. Methanogens associated with rumen ciliates. *FEMS Microbiology Ecology* 22, pp. 137-143. doi: 10.1111/j.1574-6941.1997.tb00365.x

- Ueda, Y., Onodera, R. and Kandats, M. 1975. Metabolism of peptides by rumen ciliate protozoa. *Japanese Journal of Zootechnical Science* 46, pp. 34-41.
- Ungerfeld, E. M. 2013. A theoretical comparison between two ruminal electron sinks. *Frontiers in Microbiology* 4, pp. 319-334. doi: 10.3389/fmicb.2013.00319
- Ungerfeld, E. M. 2015. Shifts in metabolic hydrogen sinks in the methanogenesis-inhibited ruminal fermentation: a meta-analysis. *Frontiers in Microbiology* 6, pp. 1-17. doi: 10.3389/fmicb.2015.00037
- Ushida, K. and Jouany, J. P. 1990. Effect of defaunation on fibre digestion in sheep given two isonitrogenous diets. *Animal Feed Science and Technology* 29, pp. 153-158. [https://doi.org/10.1016/0377-8401\(90\)90101-D](https://doi.org/10.1016/0377-8401(90)90101-D)
- Wereszka, K. and Michałowski, T. 2012. The ability of the rumen ciliate protozoan *Diploplastron affine* to digest and ferment starch. *Folia Microbiologica* 57, pp. 375-377.
- Whitelaw, F. G., Eadie, J. M., Bruce, L. A. and Shand, W. J. 1984. Methane formation in faunated and ciliate-free cattle and its relationship with rumen volatile fatty acid proportions. *British Journal of Nutrition* 52, pp. 261-275. doi:10.1079/bjn19840094
- Williams, A. G. 1986. Rumen holotrich ciliate protozoa. *Microbiological Reviews* 50, pp. 25-49.
- Williams, A. G. and Coleman, G. S. 1988. The rumen protozoa. In 'The rumen microbial ecosystem.' (Ed. PN Hobson) (ELSEVIER Applied Science: London and New York).
- Williams, A. G. 1989. Metabolic activities of rumen protozoa. In 'The role of protozoa and fungi in ruminant digestion.' (Eds. JV, Nolan, RA, Leng, DI, Demeyer) (Penambul books: Armidale, N.S.W., 2351. Australia).
- Williams, A. G. and Coleman, G. S. 1992. The rumen protozoa. (Springer-Verlag: New York, NY, USA).
- Yáñez-Ruiz, D. R., Scollan, N. D., Merry, R. J. and Newbold, C. J. 2006. Contribution of rumen protozoa to duodenal flow of nitrogen, conjugated linoleic acid and vaccenic acid in steers fed silages differing in their water-soluble carbohydrate content. *British Journal of Nutrition* 96, pp. 861-869.
- Yoder, R. D., Trenkle, A. and Burroughs, W. 1966. Influence of rumen protozoa and bacteria upon cellulose digestion in vitro. *Journal of Animal Science* 25, pp. 609-612.
- Zeitz, J. O., Amelchanka, S. L., Michałowski, T., Wereszka, K., Meile, L., Hartnack, S., Kreuzer, M. and Soliva, C. R. 2012. Effect of the rumen ciliates *Entodinium caudatum*, *Epidinium ecaudatum* and *Eudiplodinium maggii*, and combinations thereof, on ruminal fermentation and total tract digestion in sheep. *Archives of Animal Nutrition* 66, pp. 180-199. doi:10.1080/1745039x.2012.676817.

Received date: 02/02/2020

Acceptance date: 28/02/2020